

SAS/ETS[®] 14.2 User's Guide

Forecasting Process

Details

This document is an individual chapter from *SAS/ETS® 14.2 User's Guide*.

The correct bibliographic citation for this manual is as follows: SAS Institute Inc. 2016. *SAS/ETS® 14.2 User's Guide*. Cary, NC: SAS Institute Inc.

SAS/ETS® 14.2 User's Guide

Copyright © 2016, SAS Institute Inc., Cary, NC, USA

All Rights Reserved. Produced in the United States of America.

For a hard-copy book: No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without the prior written permission of the publisher, SAS Institute Inc.

For a web download or e-book: Your use of this publication shall be governed by the terms established by the vendor at the time you acquire this publication.

The scanning, uploading, and distribution of this book via the Internet or any other means without the permission of the publisher is illegal and punishable by law. Please purchase only authorized electronic editions and do not participate in or encourage electronic piracy of copyrighted materials. Your support of others' rights is appreciated.

U.S. Government License Rights; Restricted Rights: The Software and its documentation is commercial computer software developed at private expense and is provided with RESTRICTED RIGHTS to the United States Government. Use, duplication, or disclosure of the Software by the United States Government is subject to the license terms of this Agreement pursuant to, as applicable, FAR 12.212, DFAR 227.7202-1(a), DFAR 227.7202-3(a), and DFAR 227.7202-4, and, to the extent required under U.S. federal law, the minimum restricted rights as set out in FAR 52.227-19 (DEC 2007). If FAR 52.227-19 is applicable, this provision serves as notice under clause (c) thereof and no other notice is required to be affixed to the Software or documentation. The Government's rights in Software and documentation shall be only those set forth in this Agreement.

SAS Institute Inc., SAS Campus Drive, Cary, NC 27513-2414

November 2016

SAS® and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc. in the USA and other countries. ® indicates USA registration.

Other brand and product names are trademarks of their respective companies.

SAS software may be provided with certain third-party software, including but not limited to open-source software, which is licensed under its applicable third-party software license agreement. For license information about third-party software distributed with SAS software, refer to <http://support.sas.com/thirdpartylicenses>.

Chapter 63

Forecasting Process Details

Contents

Forecasting Process Summary	4035
Parameter Estimation	4036
Model Evaluation	4036
Forecasting	4038
Forecast Combination Models	4040
External or User-Supplied Forecasts	4040
Adjustments	4040
Series Transformations	4041
Smoothing Models	4042
Smoothing Model Calculations	4042
Missing Values	4044
Predictions and Prediction Errors	4044
Smoothing Weights	4044
Equations for the Smoothing Models	4046
ARIMA Models	4053
Notation for ARIMA Models	4054
Predictor Series	4057
Time Trend Curves	4057
Intervention Effects	4058
Seasonal Dummy Inputs	4060
Series Diagnostic Tests	4060
Statistics of Fit	4061
References	4063

This chapter provides computational details about several aspects of the Time Series Forecasting System.


Forecasting Process Summary

This section summarizes the forecasting process.

Parameter Estimation

The parameter estimation process for ARIMA and smoothing models is described graphically in [Figure 63.1](#).

Figure 63.1 Model Fitting Flow Diagram




The specification of smoothing and ARIMA models is described in Chapter 58, “[Specifying Forecasting Models](#).” Computational details about these kinds of models are provided in the sections “[Smoothing Models](#)” on page 4042 and “[ARIMA Models](#)” on page 4053. The results of the parameter estimation process are displayed in the Parameter Estimates table of the Model Viewer windows along with the estimate of the model variance and the final smoothing state.

Model Evaluation

The model evaluation process is described graphically in [Figure 63.2](#).


Figure 63.2 Model Evaluation Flow Diagram



Model evaluation is based on the one-step-ahead prediction errors for observations within the period of evaluation. The one-step-ahead predictions are generated from the model specification and parameter estimates. The predictions are inverse transformed (median or mean) and adjustments are removed. The prediction errors (the difference of the dependent series and the predictions) are used to compute the statistics of fit, which are described in the section “[Series Diagnostic Tests](#)” on page 4060. The results generated by the evaluation process are displayed in the Statistics of Fit table of the Model Viewer window.

Forecasting

The forecasting generation process is described graphically in [Figure 63.3](#).

Figure 63.3 Forecasting Flow Diagram

The forecasting process is similar to the model evaluation process described in the preceding section, except that k -step-ahead predictions are made from the end of the data through the specified forecast horizon, and prediction standard errors and confidence limits are calculated. The forecasts and confidence limits are displayed in the Forecast plot or table of the Model Viewer window.

Forecast Combination Models

This section discusses the computation of predicted values and confidence limits for forecast combination models. For information about how to specify forecast combination models and their combining weights, see Chapter 58, “Specifying Forecasting Models.”

Given the response time series $\{y_t : 1 \leq t \leq n\}$ with previously generated forecasts for the m component models, a combined forecast is created from the component forecasts as follows:

$$\begin{aligned} \text{Predictions:} \quad & \hat{y}_t = \sum_{i=1}^m w_i \hat{y}_{i,t} \\ \text{Prediction errors:} \quad & \hat{e}_t = y_t - \hat{y}_t \end{aligned}$$

where $\hat{y}_{i,t}$ are the forecasts of the component models and w_i are the combining weights.

The estimate of the root mean square prediction error and forecast confidence limits for the combined forecast are computed by assuming independence of the prediction errors of the component forecasts, as follows:

$$\begin{aligned} \text{Standard errors:} \quad & \hat{\sigma}_t = \sqrt{\sum_{i=1}^m w_i^2 \hat{\sigma}_{i,t}^2} \\ \text{Confidence limits:} \quad & \pm \hat{\sigma}_t Z_{\alpha/2} \end{aligned}$$

where $\hat{\sigma}_{i,t}$ are the estimated root mean square prediction errors for the component models, α is the confidence limit width, $1 - \alpha$ is the confidence level, and $Z_{\alpha/2}$ is the $\frac{\alpha}{2}$ quantile of the standard normal distribution.

Since, in practice, there might be positive correlation between the prediction errors of the component forecasts, these confidence limits may be too narrow.

External or User-Supplied Forecasts

This section discusses the computation of predicted values and confidence limits for external forecast models.

Given a response time series y_t and external forecast series $\hat{y}_t$, the prediction errors are computed as $\hat{e}_t = y_t - \hat{y}_t$ for those t for which both y_t and $\hat{y}_t$ are nonmissing. The mean squared error (MSE) is computed from the prediction errors.

The variance of the k -step-ahead prediction errors is set to k times the MSE. From these variances, the standard errors and confidence limits are computed in the usual way. If the supplied predictions contain so many missing values within the time range of the response series that the MSE estimate cannot be computed, the confidence limits, standard errors, and statistics of fit are set to missing.

Adjustments

Adjustment predictors are subtracted from the response time series prior to model parameter estimation, evaluation, and forecasting. After the predictions of the adjusted response time series are obtained from the forecasting model, the adjustments are added back to produce the forecasts.

If y_t is the response time series and $X_{i,t}$, $1 \leq i \leq m$ are m adjustment predictor series, then the adjusted response series w_t is

$$w_t = y_t - \sum_{i=1}^m X_{i,t}$$

Parameter estimation for the model is performed by using the adjusted response time series w_t . The forecasts $\hat{w}_t$ of w_t are adjusted to obtain the forecasts $\hat{y}_t$ of y_t .

$$\hat{y}_t = \hat{w}_t + \sum_{i=1}^m X_{i,t}$$

Missing values in an adjustment series are ignored in these computations.

Series Transformations

For pure ARIMA models, transforming the response time series can aid in obtaining stationary noise series. For general ARIMA models with inputs, transforming the response time series or one or more of the input time series can provide a better model fit. Similarly, the fit of smoothing models can improve when the response series is transformed.

There are four transformations available, for strictly positive series only. Let $y_t > 0$ be the original time series, and let w_t be the transformed series. The transformations are defined as follows:

Log is the logarithmic transformation,

$$w_t = \ln(y_t)$$

Logistic is the logistic transformation,

$$w_t = \ln(cy_t/(1 - cy_t))$$

where the scaling factor c is

$$c = (1 - 10^{-6})10^{-\text{ceil}(\log_{10}(\max(y_t)))}$$

and $\text{ceil}(x)$ is the smallest integer greater than or equal to x .

Square Root is the square root transformation,

$$w_t = \sqrt{y_t}$$

Box-Cox is the Box-Cox transformation,

$$w_t = \begin{cases} \frac{y_t^\lambda - 1}{\lambda}, & \lambda \neq 0 \\ \ln(y_t), & \lambda = 0 \end{cases}$$

Parameter estimation is performed by using the transformed series. The transformed model predictions and confidence limits are then obtained from the transformed time series and these parameter estimates.

The transformed model predictions $\hat{w}_t$ are used to obtain either the minimum mean absolute error (MMAE) or minimum mean squared error (MMSE) predictions $\hat{y}_t$, depending on the setting of the forecast options. The model is then evaluated based on the residuals of the original time series and these predictions. The transformed model confidence limits are inverse-transformed to obtain the forecast confidence limits.

Predictions for Transformed Models

Since the transformations described in the previous section are monotonic, applying the inverse-transformation to the transformed model predictions results in the *median* of the conditional probability density function at each point in time. This is the minimum mean absolute error (MMAE) prediction.

If $w_t = F(y_t)$ is the transform with inverse-transform $y_t = F^{-1}(w_t)$, then

$$\text{median}(\hat{y}_t) = F^{-1}(E[w_t]) = F^{-1}(\hat{w}_t)$$

The minimum mean squared error (MMSE) predictions are the *mean* of the conditional probability density function at each point in time. Assuming that the prediction errors are normally distributed with variance σ_t^2 , the MMSE predictions for each of the transformations are as follows:

Log is the conditional expectation of inverse-logarithmic transformation,

$$\hat{y}_t = E[e^{w_t}] = \exp(\hat{w}_t + \sigma_t^2/2)$$

Logistic is the conditional expectation of inverse-logistic transformation,

$$\hat{y}_t = E\left[\frac{1}{c(1 + \exp(-w_t))}\right]$$

where the scaling factor $c = (1 - e^{-6})10^{-\text{ceil}(\log_{10}(\max(y_t)))}$.

Square Root is the conditional expectation of the inverse-square root transformation,

$$\hat{y}_t = E[w_t^2] = \hat{w}_t^2 + \sigma_t^2$$

Box-Cox is the conditional expectation of the inverse Box-Cox transformation,

$$\hat{y}_t = \begin{cases} E[(\lambda w_t + 1)^{1/\lambda}], & \lambda \neq 0 \\ E[e^{w_t}] = \exp(\hat{w}_t + \frac{1}{2}\sigma_t^2), & \lambda = 0 \end{cases}$$

The expectations of the inverse logistic and Box-Cox ($\lambda \neq 0$) transformations do not generally have explicit solutions and are computed by using numerical integration.

Smoothing Models

This section details the computations performed for the exponential smoothing and Winters method forecasting models.

Smoothing Model Calculations

The descriptions and properties of various smoothing methods can be found in Gardner (1985); Chatfield (1978); Bowerman and O'Connell (1979). The following section summarizes the smoothing model computations.

Given a time series $\{Y_t : 1 \leq t \leq n\}$, the underlying model assumed by the smoothing models has the following (additive seasonal) form,

$$Y_t = \mu_t + \beta_t t + s_p(t) + \epsilon_t$$

where

μ_t	represents the time-varying mean term.
β_t	represents the time-varying slope.
$s_p(t)$	represents the time-varying seasonal contribution for one of the p seasons.
ϵ_t	are disturbances.

For smoothing models without trend terms, $\beta_t = 0$; and for smoothing models without seasonal terms, $s_p(t) = 0$. Each smoothing model is described in the following sections.

At each time t , the smoothing models estimate the time-varying components described above with the *smoothing state*. After initialization, the smoothing state is updated for each observation using the *smoothing equations*. The smoothing state at the last nonmissing observation is used for predictions.

Smoothing State and Smoothing Equations

Depending on the smoothing model, the *smoothing state* at time t consists of the following:

- L_t is a smoothed level that estimates μ_t .
- T_t is a smoothed trend that estimates β_t .
- S_{t-j} , $j = 0, \dots, p-1$, are seasonal factors that estimate $s_p(t)$.

The smoothing process starts with an initial estimate of the smoothing state, which is subsequently updated for each observation by using the *smoothing equations*.

The smoothing equations determine how the smoothing state changes as time progresses. Knowledge of the smoothing state at time $t-1$ and that of the time series value at time t uniquely determine the smoothing state at time t . The *smoothing weights* determine the contribution of the previous smoothing state to the current smoothing state. The smoothing equations for each smoothing model are listed in the following sections.

Smoothing State Initialization

Given a time series $\{Y_t : 1 \leq t \leq n\}$, the smoothing process first computes the smoothing state for time $t = 1$. However, this computation requires an initial estimate of the smoothing state at time $t = 0$, even though no data exist at or before time $t = 0$.

An appropriate choice for the initial smoothing state is made by backcasting from time $t = n$ to $t = 1$ to obtain a prediction at $t = 0$. The initialization for the backcast is obtained by regression with constant and linear terms and seasonal dummies (additive or multiplicative) as appropriate for the smoothing model. For models with linear or seasonal terms, the estimates obtained by the regression are used for initial smoothed trend and seasonal factors; however, the initial smoothed level for backcasting is always set to the last observation, Y_n .

The smoothing state at time $t = 0$ obtained from the backcast is used to initialize the smoothing process from time $t = 1$ to $t = n$ (Chatfield and Yar 1988).

For models with seasonal terms, the smoothing state is normalized so that the seasonal factors S_{t-j} for $j = 0, \dots, p - 1$ sum to zero for models that assume additive seasonality and average to one for models (such as Winters method models) that assume multiplicative seasonality.

Missing Values

When a missing value is encountered at time t , the smoothed values are updated using the *error-correction form* of the smoothing equations with the one-step-ahead prediction error, e_t , set to zero. The missing value is estimated using the one-step-ahead prediction at time $t - 1$ —that is, $\hat{Y}_{t-1}(1)$ (Aldrin and Damsleth 1989). The error-correction forms of each of the smoothing models are listed in the following sections.

Predictions and Prediction Errors

Predictions are made based on the last known smoothing state. Predictions made at time t for k steps ahead are denoted $\hat{Y}_t(k)$ and the associated prediction errors are denoted $e_t(k) = Y_{t+k} - \hat{Y}_t(k)$. The *prediction equation* for each smoothing model is listed in the following sections.

The *one-step-ahead predictions* refer to predictions made at time $t - 1$ for one time unit into the future—that is, $\hat{Y}_{t-1}(1)$. The *one-step-ahead prediction errors* are more simply denoted $e_t = e_{t-1}(1) = Y_t - \hat{Y}_{t-1}(1)$. The one-step-ahead prediction errors are also the model residuals, and the sum of squares of the one-step-ahead prediction errors is the objective function used in smoothing weight optimization.

The *variance of the prediction errors* are used to calculate the confidence limits (Sweet 1985; McKenzie 1986; Yar and Chatfield 1990; Chatfield and Yar 1991). The equations for the variance of the prediction errors for each smoothing model are listed in the following sections.

Note: $\text{var}(\epsilon_t)$ is estimated by the mean square of the one-step-ahead prediction errors.

Smoothing Weights

Depending on the smoothing model, the smoothing weights consist of the following:

α	is a level smoothing weight.
γ	is a trend smoothing weight.
δ	is a seasonal smoothing weight.
ϕ	is a trend damping weight.

Larger smoothing weights (less damping) permit the more recent data to have a greater influence on the predictions. Smaller smoothing weights (more damping) give less weight to recent data.

Specifying the Smoothing Weights

Typically the smoothing weights are chosen to be from zero to one. (This is intuitive because the weights associated with the past smoothing state and the value of current observation would normally sum to one.) However, each smoothing model (except Winters method—multiplicative version) has an ARIMA equivalent. Weights chosen to be within the ARIMA additive-invertible region will guarantee stable predictions (Archibald 1990; Gardner 1985). The ARIMA equivalent and the additive-invertible region for each smoothing model are listed in the following sections.

Optimizing the Smoothing Weights

Smoothing weights are determined so as to minimize the sum of squared, one-step-ahead prediction errors. The optimization is initialized by choosing from a predetermined grid the initial smoothing weights that result in the smallest sum of squared, one-step-ahead prediction errors. The optimization process is highly dependent on this initialization. It is possible that the optimization process will fail due to the inability to obtain stable initial values for the smoothing weights (Greene 1993; Judge et al. 1980), and it is possible for the optimization to result in a local minima.

The optimization process can result in weights to be chosen outside both the zero-to-one range and the ARIMA additive-invertible region. By restricting weight optimization to additive-invertible region, you can obtain a local minimum with stable predictions. Likewise, weight optimization can be restricted to the zero-to-one range or other ranges. It is also possible to fix certain weights to a specific value and optimize the remaining weights.

Standard Errors

The standard errors associated with the smoothing weights are calculated from the Hessian matrix of the sum of squared, one-step-ahead prediction errors with respect to the smoothing weights used in the optimization process.

Weights Near Zero or One

Sometimes the optimization process results in weights near zero or one.

For simple or double (Brown) exponential smoothing, a level weight near zero implies that simple differencing of the time series might be appropriate.

For linear (Holt) exponential smoothing, a level weight near zero implies that the smoothed trend is constant and that an ARIMA model with deterministic trend might be a more appropriate model.

For damped-trend linear exponential smoothing, a damping weight near one implies that linear (Holt) exponential smoothing might be a more appropriate model.

For Winters method and seasonal exponential smoothing, a seasonal weight near one implies that a nonseasonal model might be more appropriate and a seasonal weight near zero implies that deterministic seasonal factors might be present.

Equations for the Smoothing Models

Simple Exponential Smoothing

The model equation for simple exponential smoothing is

$$Y_t = \mu_t + \epsilon_t$$

The smoothing equation is

$$L_t = \alpha Y_t + (1 - \alpha)L_{t-1}$$

The error-correction form of the smoothing equation is

$$L_t = L_{t-1} + \alpha e_t$$

(Note: For missing values, $e_t = 0$.)

The k -step prediction equation is

$$\hat{Y}_t(k) = L_t$$

The ARIMA model equivalency to simple exponential smoothing is the ARIMA(0,1,1) model

$$(1 - B)Y_t = (1 - \theta B)\epsilon_t$$

$$\theta = 1 - \alpha$$

The moving-average form of the equation is

$$Y_t = \epsilon_t + \sum_{j=1}^{\infty} \alpha \epsilon_{t-j}$$

For simple exponential smoothing, the additive-invertible region is

$$\{0 < \alpha < 2\}$$

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[1 + \sum_{j=1}^{k-1} \alpha^2 \right] = \text{var}(\epsilon_t)(1 + (k-1)\alpha^2)$$

Double (Brown) Exponential Smoothing

The model equation for double exponential smoothing is

$$Y_t = \mu_t + \beta_t t + \epsilon_t$$

The smoothing equations are

$$\begin{aligned} L_t &= \alpha Y_t + (1 - \alpha) L_{t-1} \\ T_t &= \alpha (L_t - L_{t-1}) + (1 - \alpha) T_{t-1} \end{aligned}$$

This method can be equivalently described in terms of two successive applications of simple exponential smoothing,

$$\begin{aligned} S_t^{[1]} &= \alpha Y_t + (1 - \alpha) S_{t-1}^{[1]} \\ S_t^{[2]} &= \alpha S_t^{[1]} + (1 - \alpha) S_{t-1}^{[2]} \end{aligned}$$

where $S_t^{[1]}$ are the smoothed values of Y_t , and $S_t^{[2]}$ are the smoothed values of $S_t^{[1]}$. The prediction equation then takes the following form:

$$\hat{Y}_t(k) = (2 + \alpha k / (1 - \alpha)) S_t^{[1]} - (1 + \alpha k / (1 - \alpha)) S_t^{[2]}$$

The error-correction forms of the smoothing equations are

$$\begin{aligned} L_t &= L_{t-1} + T_{t-1} + \alpha e_t \\ T_t &= T_{t-1} + \alpha^2 e_t \end{aligned}$$

(Note: For missing values, $e_t = 0$.)

The k -step prediction equation is

$$\hat{Y}_t(k) = L_t + ((k - 1) + 1/\alpha) T_t$$

The ARIMA model equivalency to double exponential smoothing is the ARIMA(0,2,2) model,

$$\begin{aligned} (1 - B)^2 Y_t &= (1 - \theta B)^2 \epsilon_t \\ \theta &= 1 - \alpha \end{aligned}$$

The moving-average form of the equation is

$$Y_t = \epsilon_t + \sum_{j=1}^{\infty} (2\alpha + (j - 1)\alpha^2) \epsilon_{t-j}$$

For double exponential smoothing, the additive-invertible region is

$$\{0 < \alpha < 2\}$$

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[1 + \sum_{j=1}^{k-1} (2\alpha + (j - 1)\alpha^2)^2 \right]$$

Linear (Holt) Exponential Smoothing

The model equation for linear exponential smoothing is

$$Y_t = \mu_t + \beta_t t + \epsilon_t$$

The smoothing equations are

$$\begin{aligned} L_t &= \alpha Y_t + (1 - \alpha)(L_{t-1} + T_{t-1}) \\ T_t &= \gamma(L_t - L_{t-1}) + (1 - \gamma)T_{t-1} \end{aligned}$$

The error-correction form of the smoothing equations is

$$\begin{aligned} L_t &= L_{t-1} + T_{t-1} + \alpha e_t \\ T_t &= T_{t-1} + \alpha \gamma e_t \end{aligned}$$

(Note: For missing values, $e_t = 0$.)

The k -step prediction equation is

$$\hat{Y}_t(k) = L_t + kT_t$$

The ARIMA model equivalency to linear exponential smoothing is the ARIMA(0,2,2) model,

$$\begin{aligned} (1 - B)^2 Y_t &= (1 - \theta_1 B - \theta_2 B^2) \epsilon_t \\ \theta_1 &= 2 - \alpha - \alpha \gamma \\ \theta_2 &= \alpha - 1 \end{aligned}$$

The moving-average form of the equation is

$$Y_t = \epsilon_t + \sum_{j=1}^{\infty} (\alpha + j\alpha\gamma) \epsilon_{t-j}$$

For linear exponential smoothing, the additive-invertible region is

$$\begin{aligned} \{0 < \alpha < 2\} \\ \{0 < \gamma < 4/\alpha - 2\} \end{aligned}$$

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[1 + \sum_{j=1}^{k-1} (\alpha + j\alpha\gamma)^2 \right]$$

Damped-Trend Linear Exponential Smoothing

The model equation for damped-trend linear exponential smoothing is

$$Y_t = \mu_t + \beta_t t + \epsilon_t$$

The smoothing equations are

$$\begin{aligned} L_t &= \alpha Y_t + (1 - \alpha)(L_{t-1} + \phi T_{t-1}) \\ T_t &= \gamma(L_t - L_{t-1}) + (1 - \gamma)\phi T_{t-1} \end{aligned}$$

The error-correction form of the smoothing equations is

$$L_t = L_{t-1} + \phi T_{t-1} + \alpha e_t \quad T_t = \phi T_{t-1} + \alpha \gamma e_t$$

(Note: For missing values, $e_t = 0$.)

The k -step prediction equation is

$$\hat{Y}_t(k) = L_t + \sum_{i=1}^k \phi^i T_t$$

The ARIMA model equivalency to damped-trend linear exponential smoothing is the ARIMA(1,1,2) model,

$$\begin{aligned} (1 - \phi B)(1 - B)Y_t &= (1 - \theta_1 B - \theta_2 B^2)\epsilon_t \\ \theta_1 &= 1 + \phi - \alpha - \alpha\gamma\phi \\ \theta_2 &= (\alpha - 1)\phi \end{aligned}$$

The moving-average form of the equation (assuming $|\phi| < 1$) is

$$Y_t = \epsilon_t + \sum_{j=1}^{\infty} (\alpha + \alpha\gamma\phi(\phi^j - 1)/(\phi - 1))\epsilon_{t-j}$$

For damped-trend linear exponential smoothing, the additive-invertible region is

$$\begin{aligned} \{0 < \alpha < 2\} \\ \{0 < \phi\gamma < 4/\alpha - 2\} \end{aligned}$$

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[1 + \sum_{j=1}^{k-1} (\alpha + \alpha\gamma\phi(\phi^j - 1)/(\phi - 1))^2 \right]$$

Seasonal Exponential Smoothing

The model equation for seasonal exponential smoothing is

$$Y_t = \mu_t + s_p(t) + \epsilon_t$$

The smoothing equations are

$$\begin{aligned} L_t &= \alpha(Y_t - S_{t-p}) + (1 - \alpha)L_{t-1} \\ S_t &= \delta(Y_t - L_t) + (1 - \delta)S_{t-p} \end{aligned}$$

The error-correction form of the smoothing equations is

$$\begin{aligned} L_t &= L_{t-1} + \alpha e_t \\ S_t &= S_{t-p} + \delta(1 - \alpha)e_t \end{aligned}$$

(Note: For missing values, $e_t = 0$.)

The k -step prediction equation is

$$\hat{Y}_t(k) = L_t + S_{t-p+k}$$

The ARIMA model equivalency to seasonal exponential smoothing is the $\text{ARIMA}(0,1,p+1)(0,1,0)_p$ model,

$$\begin{aligned} (1 - B)(1 - B^p)Y_t &= (1 - \theta_1 B - \theta_2 B^p - \theta_3 B^{p+1})\epsilon_t \\ \theta_1 &= 1 - \alpha \\ \theta_2 &= 1 - \delta(1 - \alpha) \\ \theta_3 &= (1 - \alpha)(\delta - 1) \end{aligned}$$

The moving-average form of the equation is

$$\begin{aligned} Y_t &= \epsilon_t + \sum_{j=1}^{\infty} \psi_j \epsilon_{t-j} \\ \psi_j &= \begin{cases} \alpha & \text{for } j \bmod p \neq 0 \\ \alpha + \delta(1 - \alpha) & \text{for } j \bmod p = 0 \end{cases} \end{aligned}$$

For seasonal exponential smoothing, the additive-invertible region is

$$\{\max(-p\alpha, 0) < \delta(1 - \alpha) < (2 - \alpha)\}$$

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[1 + \sum_{j=1}^{k-1} \psi_j^2 \right]$$

Multiplicative Seasonal Smoothing

In order to use the multiplicative version of seasonal smoothing, the time series and all predictions must be strictly positive.

The model equation for the multiplicative version of seasonal smoothing is

$$Y_t = \mu_t s_p(t) + \epsilon_t$$

The smoothing equations are

$$\begin{aligned} L_t &= \alpha(Y_t/S_{t-p}) + (1 - \alpha)L_{t-1} \\ S_t &= \delta(Y_t/L_t) + (1 - \delta)S_{t-p} \end{aligned}$$

The error-correction form of the smoothing equations is

$$\begin{aligned} L_t &= L_{t-1} + \alpha e_t / S_{t-p} \\ S_t &= S_{t-p} + \delta(1 - \alpha)e_t / L_t \end{aligned}$$

(Note: For missing values, $e_t = 0$.)

The k -step prediction equation is

$$\hat{Y}_t(k) = L_t S_{t-p+k}$$

The multiplicative version of seasonal smoothing does not have an ARIMA equivalent; however, when the seasonal variation is small, the ARIMA additive-invertible region of the additive version of seasonal described in the preceding section can approximate the stability region of the multiplicative version.

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[\sum_{i=0}^{\infty} \sum_{j=0}^{p-1} (\psi_{j+ip} S_{t+k} / S_{t+k-j})^2 \right]$$

where ψ_j are as described for the additive version of seasonal method, and $\psi_j = 0$ for $j \geq k$.

Winters Method—Additive Version

The model equation for the additive version of the Winters method is

$$Y_t = \mu_t + \beta_t t + s_p(t) + \epsilon_t$$

The smoothing equations are

$$\begin{aligned} L_t &= \alpha(Y_t - S_{t-p}) + (1 - \alpha)(L_{t-1} + T_{t-1}) \\ T_t &= \gamma(L_t - L_{t-1}) + (1 - \gamma)T_{t-1} \\ S_t &= \delta(Y_t - L_t) + (1 - \delta)S_{t-p} \end{aligned}$$

The error-correction form of the smoothing equations is

$$\begin{aligned} L_t &= L_{t-1} + T_{t-1} + \alpha e_t \\ T_t &= T_{t-1} + \alpha \gamma e_t \\ S_t &= S_{t-p} + \delta(1 - \alpha)e_t \end{aligned}$$

(Note: For missing values, $e_t = 0$.)

The k -step prediction equation is

$$\hat{Y}_t(k) = L_t + kT_t + S_{t-p+k}$$

The ARIMA model equivalency to the additive version of the Winters method is the $\text{ARIMA}(0,1,p+1)(0,1,0)_p$ model,

$$\begin{aligned} (1 - B)(1 - B^p)Y_t &= \left[1 - \sum_{i=1}^{p+1} \theta_i B^i \right] \epsilon_t \\ \theta_j &= \begin{cases} 1 - \alpha - \alpha\gamma & j = 1 \\ -\alpha\gamma & 2 \leq j \leq p-1 \\ 1 - \alpha\gamma - \delta(1 - \alpha) & j = p \\ (1 - \alpha)(\delta - 1) & j = p+1 \end{cases} \end{aligned}$$

The moving-average form of the equation is

$$\begin{aligned} Y_t &= \epsilon_t + \sum_{j=1}^{\infty} \psi_j \epsilon_{t-j} \\ \psi_j &= \begin{cases} \alpha + j\alpha\gamma & \text{for } j \bmod p \neq 0 \\ \alpha + j\alpha\gamma + \delta(1 - \alpha), & \text{for } j \bmod p = 0 \end{cases} \end{aligned}$$

For the additive version of the Winters method (Archibald 1990), the additive-invertible region is

$$\begin{aligned} &\{\max(-p\alpha, 0) < \delta(1 - \alpha) < (2 - \alpha)\} \\ &\{0 < \alpha\gamma < 2 - \alpha - \delta(1 - \alpha)(1 - \cos(\vartheta))\} \end{aligned}$$

where ϑ is the smallest nonnegative solution to the equations listed in Archibald (1990).

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[1 + \sum_{j=1}^{k-1} \psi_j^2 \right]$$

Winters Method—Multiplicative Version

In order to use the multiplicative version of the Winters method, the time series and all predictions must be strictly positive.

The model equation for the multiplicative version of the Winters method is

$$Y_t = (\mu_t + \beta_t t)s_p(t) + \epsilon_t$$

The smoothing equations are

$$L_t = \alpha(Y_t/S_{t-p}) + (1 - \alpha)(L_{t-1} + T_{t-1})$$

$$T_t = \gamma(L_t - L_{t-1}) + (1 - \gamma)T_{t-1}$$

$$S_t = \delta(Y_t/L_t) + (1 - \delta)S_{t-p}$$

The error-correction form of the smoothing equations is

$$L_t = L_{t-1} + T_{t-1} + \alpha e_t/S_{t-p}$$

$$T_t = T_{t-1} + \alpha \gamma e_t/S_{t-p}$$

$$S_t = S_{t-p} + \delta(1 - \alpha)e_t/L_t$$

NOTE: For missing values, $e_t = 0$.

The k -step prediction equation is

$$\hat{Y}_t(k) = (L_t + kT_t)S_{t-p+k}$$

The multiplicative version of the Winters method does not have an ARIMA equivalent; however, when the seasonal variation is small, the ARIMA additive-invertible region of the additive version of the Winters method described in the preceding section can approximate the stability region of the multiplicative version.

The variance of the prediction errors is estimated as

$$\text{var}(e_t(k)) = \text{var}(\epsilon_t) \left[\sum_{i=0}^{\infty} \sum_{j=0}^{p-1} (\psi_{j+ip} S_{t+k}/S_{t+k-j})^2 \right]$$

where ψ_j are as described for the additive version of the Winters method and $\psi_j = 0$ for $j \geq k$.

ARIMA Models

Autoregressive integrated moving-average (ARIMA) models predict values of a dependent time series with a linear combination of its own past values, past errors (also called shocks or innovations), and current and past values of other time series (predictor time series).

The Time Series Forecasting System uses the ARIMA procedure of SAS/ETS software to fit and forecast ARIMA models. The maximum likelihood method is used for parameter estimation. For more information about ARIMA model estimation and forecasting, see Chapter 7, “[The ARIMA Procedure](#).”

This section summarizes the notation used for ARIMA models.

Notation for ARIMA Models

A dependent time series that is modeled as a linear combination of its own past values and past values of an error series is known as a (pure) ARIMA model.

Nonseasonal ARIMA Model Notation

The order of an ARIMA model is usually denoted by the notation $\text{ARIMA}(p,d,q)$, where

p	is the order of the autoregressive part.
d	is the order of the differencing (rarely should $d > 2$ be needed).
q	is the order of the moving-average process.

Given a dependent time series $\{Y_t : 1 \leq t \leq n\}$, mathematically the ARIMA model is written as

$$(1 - B)^d Y_t = \mu + \frac{\theta(B)}{\phi(B)} a_t$$

where

t	indexes time.
μ	is the mean term.
B	is the backshift operator; that is, $BX_t = X_{t-1}$.
$\phi(B)$	is the autoregressive operator, represented as a polynomial in the backshift operator: $\phi(B) = 1 - \phi_1 B - \dots - \phi_p B^p$.
$\theta(B)$	is the moving-average operator, represented as a polynomial in the backshift operator: $\theta(B) = 1 - \theta_1 B - \dots - \theta_q B^q$.
a_t	is the independent disturbance, also called the random error.

For example, the mathematical form of the $\text{ARIMA}(1,1,2)$ model is

$$(1 - B)Y_t = \mu + \frac{(1 - \theta_1 B - \theta_2 B^2)}{(1 - \phi_1 B)} a_t$$

Seasonal ARIMA Model Notation

Seasonal ARIMA models are expressed in factored form by the notation $\text{ARIMA}(p,d,q)(P,D,Q)_s$, where

P	is the order of the seasonal autoregressive part.
D	is the order of the seasonal differencing (rarely should $D > 1$ be needed).
Q	is the order of the seasonal moving-average process.
s	is the length of the seasonal cycle.

Given a dependent time series $\{Y_t : 1 \leq t \leq n\}$, mathematically the ARIMA seasonal model is written as

$$(1 - B)^d (1 - B^s)^D Y_t = \mu + \frac{\theta(B)\theta_s(B^s)}{\phi(B)\phi_s(B^s)} a_t$$

where

$\phi_s(B^s)$	is the seasonal autoregressive operator, represented as a polynomial in the backshift operator: $\phi_s(B^s) = 1 - \phi_{s,1}B^s - \cdots - \phi_{s,P}B^{sP}$.
$\theta_s(B^s)$	is the seasonal moving-average operator, represented as a polynomial in the backshift operator: $\theta_s(B^s) = 1 - \theta_{s,1}B^s - \cdots - \theta_{s,Q}B^{sQ}$.

For example, the mathematical form of the ARIMA(1,0,1)(1,1,2)₁₂ model is

$$(1 - B^{12})Y_t = \mu + \frac{(1 - \theta_1 B)(1 - \theta_{s,1}B^{12} - \theta_{s,2}B^{24})}{(1 - \phi_1 B)(1 - \phi_{s,1}B^{12})}a_t$$

Abbreviated Notation for ARIMA Models

If the differencing order, autoregressive order, or moving-average order is zero, the notation is further abbreviated as follows:

$I(d)(D)_s$	integrated model or ARIMA(0,d,0)(0,D,0)
$AR(p)(P)_s$	autoregressive model or ARIMA(p,0,0)(P,0,0)
$IAR(p,d)(P,D)_s$	integrated autoregressive model or ARIMA(p,d,0)(P,D,0) _s
$MA(q)(Q)_s$	moving average model or ARIMA(0,0,q)(0,0,Q) _s
$IMA(d,q)(D,Q)_s$	integrated moving average model or ARIMA(0,d,q)(0,D,Q) _s
$ARMA(p,q)(P,Q)_s$	autoregressive moving-average model or ARIMA(p,0,q)(P,0,Q) _s

Notation for Transfer Functions

A transfer function can be used to filter a predictor time series to form a dynamic regression model.

Let Y_t be the dependent series, let X_t be the predictor series, and let $\Psi(B)$ be a linear filter or transfer function for the effect of X_t on Y_t . The ARIMA model is then

$$(1 - B)^d(1 - B^s)^D Y_t = \mu + \Psi(B)(1 - B)^d(1 - B^s)^D X_t + \frac{\theta(B)\theta_s(B^s)}{\phi(B)\phi_s(B^s)}a_t$$

This model is called a *dynamic regression* of Y_t on X_t .

Nonseasonal Transfer Function Notation

Given the i th predictor time series $\{X_{i,t} : 1 \leq t \leq n\}$, the transfer function is written as

$$\text{Dif}(d_i)\text{Lag}(k_i)\text{N}(q_i)/\text{D}(p_i)$$

where

d_i	is the simple order of the differencing for the i th predictor time series, $(1 - B)^{d_i}X_{i,t}$ (rarely should $d_i > 2$ be needed).
k_i	is the pure time delay (lag) for the effect of the i th predictor time series, $X_{i,t}B^{k_i} = X_{i,t-k_i}$.

p_i	is the simple order of the denominator for the i th predictor time series.
q_i	is the simple order of the numerator for the i th predictor time series.

The mathematical notation used to describe a transfer function is

$$\Psi_i(B) = \frac{\omega_i(B)}{\delta_i(B)} (1 - B)^{d_i} B^{k_i}$$

where

B	is the backshift operator; that is, $BX_t = X_{t-1}$.
$\delta_i(B)$	is the denominator polynomial of the transfer function for the i th predictor time series: $\delta_i(B) = 1 - \delta_{i,1}B - \dots - \delta_{i,p_i}B^{p_i}$.
$\omega_i(B)$	is the numerator polynomial of the transfer function for the i th predictor time series: $\omega_i(B) = 1 - \omega_{i,1}B - \dots - \omega_{i,q_i}B^{q_i}$.

The numerator factors for a transfer function for a predictor series are like the MA part of the ARMA model for the noise series. The denominator factors for a transfer function for a predictor series are like the AR part of the ARMA model for the noise series. Denominator factors introduce exponentially weighted, infinite distributed lags into the transfer function.

For example, the transfer function for the i th predictor time series with

$k_i = 3$	time lag is 3
$d_i = 1$	simple order of differencing is one
$p_i = 1$	simple order of the denominator is one
$q_i = 2$	simple order of the numerator is two

would be written as [Dif(1)Lag(3)N(2)/D(1)]. The mathematical notation for the transfer function in this example is

$$\Psi_i(B) = \frac{(1 - \omega_{i,1}B - \omega_{i,2}B^2)}{(1 - \delta_{i,1}B)} (1 - B)B^3$$

Seasonal Transfer Function Notation

The general transfer function notation for the i th predictor time series $X_{i,t}$ with seasonal factors is [Dif(d_i)(D_i)_s Lag(k_i) N(q_i)(Q_i)_s/ D(p_i)(P_i)_s], where

D_i	is the seasonal order of the differencing for the i th predictor time series (rarely should $D_i > 1$ be needed).
P_i	is the seasonal order of the denominator for the i th predictor time series (rarely should $P_i > 2$ be needed).
Q_i	is the seasonal order of the numerator for the i th predictor time series, (rarely should $Q_i > 2$ be needed).
s	is the length of the seasonal cycle.

The mathematical notation used to describe a seasonal transfer function is

$$\Psi_i(B) = \frac{\omega_i(B)\omega_{s,i}(B^s)}{\delta_i(B)\delta_{s,i}(B^s)}(1-B)^{d_i}(1-B^s)^{D_i}B^{k_i}$$

where

$\delta_{s,i}(B^s)$ is the denominator seasonal polynomial of the transfer function for the i th predictor time series: $\delta_{s,i}(B) = 1 - \delta_{s,i,1}B - \dots - \delta_{s,i,P_i}B^{sP_i}$.

$\omega_{s,i}(B^s)$ is the numerator seasonal polynomial of the transfer function for the i th predictor time series: $\omega_{s,i}(B) = 1 - \omega_{s,i,1}B - \dots - \omega_{s,i,Q_i}B^{sQ_i}$.

For example, the transfer function for the i th predictor time series $X_{i,t}$ whose seasonal cycle $s = 12$ with

$d_i = 2$ simple order of differencing is two
 $D_i = 1$ seasonal order of differencing is one
 $q_i = 2$ simple order of the numerator is two
 $Q_i = 1$ seasonal order of the numerator is one

would be written as $[\text{Dif}(2)(1)_s \text{N}(2)(1)_s]$. The mathematical notation for the transfer function in this example is

$$\Psi_i(B) = (1 - \omega_{i,1}B - \omega_{i,2}B^2)(1 - \omega_{s,i,1}B^{12})(1-B)^2(1-B^{12})$$

Note: In this case, $[\text{Dif}(2)(1)_s \text{N}(2)(1)_s] = [\text{Dif}(2)(1)_s \text{Lag}(0)\text{N}(2)(1)_s/\text{D}(0)(0)_s]$.

Predictor Series

This section discusses time trend curves, seasonal dummies, interventions, and adjustments.

Time Trend Curves

When you specify a time trend curve as a predictor in a forecasting model, the system computes a predictor series that is a deterministic function of time. This variable is then included in the model as a regressor, and the trend curve is fit to the dependent series by linear regression, in addition to other predictor series.

Some kinds of nonlinear trend curves are fit by transforming the dependent series. For example, the exponential trend curve is actually a linear time trend fit to the logarithm of the series. For these trend curve specifications, the series transformation option is set automatically, and you cannot independently control both the time trend curve and transformation option.

The computed time trend variable is included in the output data set in a variable named in accordance with the trend curve type. Let t represent the observation count from the start of the period of fit for the model, and let X_t represent the value of the time trend variable at observation t within the period of fit. The names and definitions of these variables are as follows. (Note: These deterministic variables are reserved variable names.)

Linear trend	variable name <code>_LINEAR_</code> , with $X_t = t - c$
Quadratic trend	variable name <code>_QUAD_</code> , with $X_t = (t - c)^2$. Note that a quadratic trend implies a linear trend as a special case and results in two regressors: <code>_QUAD_</code> and <code>_LINEAR_</code> .
Cubic trend	variable name <code>_CUBE_</code> , with $X_t = (t - c)^3$. Note that a cubic trend implies a quadratic trend as a special case and results in three regressors: <code>_CUBE_</code> , <code>_QUAD_</code> , and <code>_LINEAR_</code> .
Logistic trend	variable name <code>_LOGIT_</code> , with $X_t = t$. The model is a linear time trend applied to the logistic transform of the dependent series. Thus, specifying a logistic trend is equivalent to specifying the logistic series transformation and a linear time trend. A logistic trend predictor can be used only in conjunction with the logistic transformation, which is set automatically when you specify logistic trend.
Logarithmic trend	variable name <code>_LOG_</code> , with $X_t = \ln(t)$
Exponential trend	variable name <code>_EXP_</code> , with $X_t = t$. The model is a linear time trend applied to the logarithms of the dependent series. Thus, specifying an exponential trend is equivalent to specifying the log series transformation and a linear time trend. An exponential trend predictor can be used only in conjunction with the log transformation, which is set automatically when you specify exponential trend.
Hyperbolic trend	variable name <code>_HYP_</code> , with $X_t = 1/t$
Power curve trend	variable name <code>_POW_</code> , with $X_t = \ln(t)$. The model is a logarithmic time trend applied to the logarithms of the dependent series. Thus, specifying a power curve is equivalent to specifying the log series transformation and a logarithmic time trend. A power curve predictor can be used only in conjunction with the log transformation, which is set automatically when you specify a power curve trend.
EXP(A+B/TIME) trend	variable name <code>_ERT_</code> , with $X_t = 1/t$. The model is a hyperbolic time trend applied to the logarithms of the dependent series. Thus, specifying this trend curve is equivalent to specifying the log series transformation and a hyperbolic time trend. This trend curve can be used only in conjunction with the log transformation, which is set automatically when you specify this trend.

Intervention Effects

Interventions are used for modeling events that occur at specific times. That is, they are known changes that affect the dependent series or outliers.

The i th intervention series is included in the output data set with variable name `_INTV $i$ _`, which is a reserved variable name.

Point Interventions

The point intervention is a one-time event. The i th intervention series $X_{i,t}$ has a point intervention at time t_{int} when the series is nonzero only at time t_{int} —that is,

$$X_{i,t} = \begin{cases} 1, & t = t_{int} \\ 0, & \text{otherwise} \end{cases}$$

Step Interventions

Step interventions are continuing, and the input time series flags periods after the intervention. For a step intervention, before time t_{int} , the i th intervention series $X_{i,t}$ is zero and then steps to a constant level thereafter—that is,

$$X_{i,t} = \begin{cases} 1, & t \geq t_{int} \\ 0, & \text{otherwise} \end{cases}$$

Ramp Interventions

A ramp intervention is a continuing intervention that increases linearly after the intervention time. For a ramp intervention, before time t_{int} , the i th intervention series $X_{i,t}$ is zero and increases linearly thereafter—that is, proportional to time.

$$X_{i,t} = \begin{cases} t - t_{int}, & t \geq t_{int} \\ 0, & \text{otherwise} \end{cases}$$

Intervention Effect

Given the i th intervention series $X_{i,t}$, you can define how the intervention takes effect by filters (transfer functions) of the form

$$\Psi_i(B) = \frac{1 - \omega_{i,1}B - \dots - \omega_{i,q_i}B^{q_i}}{1 - \delta_{i,1}B - \dots - \delta_{i,p_i}B^{p_i}}$$

where B is the backshift operator $By_t = y_{t-1}$.

The denominator of the transfer function determines the decay pattern of the intervention effect, whereas the numerator terms determine the size of the intervention effect time window.

For example, the following intervention effects are associated with the respective transfer functions:

Immediately	$\Psi_i(B) = 1$
Gradually	$\Psi_i(B) = 1/(1 - \delta_{i,1}B)$
1 lag window	$\Psi_i(B) = 1 - \omega_{i,1}B$
3 lag window	$\Psi_i(B) = 1 - \omega_{i,1}B - \omega_{i,2}B^2 - \omega_{i,3}B^3$

Intervention Notation

The notation used to describe intervention effects has the form *type* : t_{int} (q_i)/(p_i), where *type* is point, step, or ramp; t_{int} is the time of the intervention (for example, OCT87); q_i is the transfer function numerator order; and p_i is the transfer function denominator order. If $q_i = 0$, the part “(q_i)” is omitted; if $p_i = 0$, the part “/(p_i)” is omitted.

In the Intervention Specification window, the `Number of Lags` option specifies the transfer function numerator order q_i , and the `Effect Decay Pattern` option specifies the transfer function denominator order p_i . In the `Effect Decay Pattern` options, values and resulting p_i are as follows: `None`, $p_i = 0$ `Exp`, $p_i = 1$ `Wave`, $p_i = 2$

For example, a step intervention with date 08MAR90 and effect pattern `Exp` is denoted “Step:08MAR90/(1)” and has a transfer function filter $\Psi_i(B) = 1/(1 - \delta_1 B)$. A ramp intervention immediately applied on 08MAR90 is denoted “Ramp:08MAR90” and has a transfer function filter $\Psi_i(B) = 1$.

Seasonal Dummy Inputs

For a seasonal cycle of length s , the seasonal dummy regressors include

$$\{X_{i,t} : 1 \leq i \leq (s-1), 1 \leq t \leq n\}$$

for models that include an intercept term and

$$\{X_{i,t} : 1 \leq i \leq s, 1 \leq t \leq n\}$$

for models that exclude an intercept term. Each element of a seasonal dummy regressor is either zero or one, based on the following rule:

$$X_{i,t} = \begin{cases} 1, & \text{when } i = t \bmod s \\ 0, & \text{otherwise} \end{cases}$$

Note that if the model includes an intercept term, the number of seasonal dummy regressors is one less than s to ensure that the linear system is full rank.

The seasonal dummy variables are included in the output data set with variable names prefixed with “SDUMMY” and sequentially numbered. They are reserved variable names.

Series Diagnostic Tests

This section describes the diagnostic tests that are used to determine the kinds of forecasting models appropriate for a series.

The series diagnostics are a set of heuristics that provide recommendations on whether or not the forecasting model should contain a log transform, trend terms, and seasonal terms. These recommendations are used by the automatic model selection process to restrict the model search to a subset of the model selection list. (You can disable this behavior by using the Automatic Model Selection Options window.)

The tests that are used by the series diagnostics do not always produce the correct classification of the series. They are intended to accelerate the process of searching for a good forecasting model for the series, but you should not rely on them if finding the very best model is important to you.

If you have information about the appropriate kinds of forecasting models (perhaps from studying the plots and autocorrelations shown in the Series Viewer window), you can set the series diagnostic flags in the Series Diagnostics window. Select the YES, NO, or MAYBE values for the `Log Transform`, `Trend`, and `Seasonality` options in the Series Diagnostics window as you think appropriate.

The series diagnostics tests are intended as a heuristic tool only, and no statistical validity is claimed for them. These tests might be modified and enhanced in future releases of the Time Series Forecasting System. The testing strategy is as follows:

1. **Log transform test.** The log test fits a high-order autoregressive model to the series and to the log of the series and compares goodness-of-fit measures for the prediction errors of the two models. If this test finds that log transforming the series is suitable, the `Log Transform` option is set to YES, and the subsequent diagnostic tests are performed on the log transformed series.

2. **Trend test.** The resultant series is tested for presence of a trend by using an augmented Dickey-Fuller test and a random walk with drift test. If either test finds that the series appears to have a trend, the `Trend` option is set to YES, and the subsequent diagnostic tests are performed on the differenced series.
3. **Seasonality test.** The resultant series is tested for seasonality. A seasonal dummy model with AR(1) errors is fit and the joint significance of the seasonal dummy estimates is tested. If the seasonal dummies are significant, the AIC statistic for this model is compared to the AIC for an AR(1) model without seasonal dummies. If the AIC for the seasonal model is lower than that of the nonseasonal model, the `Seasonal` option is set to YES.

Statistics of Fit

This section explains the goodness-of-fit statistics reported to measure how well different models fit the data. The statistics of fit for the various forecasting models can be viewed or stored in a data set by using the Model Viewer window.

Statistics of fit are computed by using the actual and forecasted values for observations in the period of evaluation. One-step forecasted values are used whenever possible, including the case when a hold-out sample contains no missing values. If a one-step forecast for an observation cannot be computed due to missing values for previous series observations, a multi-step forecast is computed, using the minimum number of steps as the previous nonmissing values in the data range permit.

The various statistics of fit reported are as follows. In these formulas, n is the number of nonmissing observations and k is the number of fitted parameters in the model.

Number of Nonmissing Observations

The number of nonmissing observations used to fit the model.

Number of Observations

The total number of observations used to fit the model, including both missing and nonmissing observations.

Number of Missing Actuals

The number of missing actual values.

Number of Missing Predicted Values

The number of missing predicted values.

Number of Model Parameters

The number of parameters fit to the data. For combined forecast, this is the number of forecast components.

Total Sum of Squares (Uncorrected)

The total sum of squares for the series, SST, uncorrected for the mean: $\sum_{t=1}^n y_t^2$.

Total Sum of Squares (Corrected)

The total sum of squares for the series, SST, corrected for the mean: $\sum_{t=1}^n (y_t - \bar{y})^2$, where $\bar{y}$ is the series mean.

Sum of Square Errors

The sum of the squared prediction errors, SSE. $SSE = \sum_{t=1}^n (y_t - \hat{y}_t)^2$, where $\hat{y}$ is the one-step predicted value.

Mean Squared Error

The mean squared prediction error, MSE, calculated from the one-step-ahead forecasts. $MSE = \frac{1}{n} SSE$. This formula enables you to evaluate small hold-out samples.

Root Mean Squared Error

The root mean square error (RMSE), $\sqrt{MSE}$.

Mean Absolute Percent Error

The mean absolute percent prediction error (MAPE), $\frac{100}{n} \sum_{t=1}^n |(y_t - \hat{y}_t)/y_t|$. The summation ignores observations where $y_t = 0$.

Mean Absolute Error

The mean absolute prediction error, $\frac{1}{n} \sum_{t=1}^n |y_t - \hat{y}_t|$.

R-Square

The R^2 statistic, $R^2 = 1 - SSE/SST$. If the model fits the series badly, the model error sum of squares, SSE, can be larger than SST and the R^2 statistic will be negative.

Adjusted R-Square

The adjusted R^2 statistic, $1 - (\frac{n-1}{n-k})(1 - R^2)$.

Amemiya's Adjusted R-Square

Amemiya's adjusted R^2 , $1 - (\frac{n+k}{n-k})(1 - R^2)$.

Random Walk R-Square

The random walk R^2 statistic (Harvey's R^2 statistic by using the random walk model for comparison), $1 - (\frac{n-1}{n})SSE/RWSSE$, where $RWSSE = \sum_{t=2}^n (y_t - y_{t-1} - \mu)^2$, and $\mu = \frac{1}{n-1} \sum_{t=2}^n (y_t - y_{t-1})$.

Akaike's Information Criterion

Akaike's information criterion (AIC), $n \ln(MSE) + 2k$.

Schwarz Bayesian Information Criterion

Schwarz Bayesian information criterion (SBC or BIC), $n \ln(MSE) + k \ln(n)$.

Amemiya's Prediction Criterion

Amemiya's prediction criterion, $\frac{1}{n} SST(\frac{n+k}{n-k})(1 - R^2) = (\frac{n+k}{n-k})\frac{1}{n} SSE$.

Maximum Error

The largest prediction error.

Minimum Error

The smallest prediction error.

Maximum Percent Error

The largest percent prediction error, $100 \max((y_t - \hat{y}_t)/y_t)$. The summation ignores observations where $y_t = 0$.

Minimum Percent Error

The smallest percent prediction error, $100 \min((y_t - \hat{y}_t)/y_t)$. The summation ignores observations where $y_t = 0$.

Mean Error

The mean prediction error, $\frac{1}{n} \sum_{t=1}^n (y_t - \hat{y}_t)$.

Mean Percent Error

The mean percent prediction error, $\frac{100}{n} \sum_{t=1}^n \frac{(y_t - \hat{y}_t)}{y_t}$. The summation ignores observations where $y_t = 0$.

References

- Akaike, H. (1974). "A New Look at the Statistical Model Identification." *IEEE Transactions on Automatic Control* AC-19:716–723.
- Aldrin, M., and Damsleth, E. (1989). "Forecasting Non-seasonal Time Series with Missing Observations." *Journal of Forecasting* 8:97–116.
- Anderson, T. W. (1971). *The Statistical Analysis of Time Series*. New York: John Wiley & Sons.
- Ansley, C. F. (1979). "An Algorithm for the Exact Likelihood of a Mixed Autoregressive–Moving Average Process." *Biometrika* 66:59–65.
- Ansley, C. F., and Newbold, P. (1980). "Finite Sample Properties of Estimators for Autoregressive Moving-Average Models." *Journal of Econometrics* 13:159–183.
- Archibald, B. C. (1990). "Parameter Space of the Holt-Winters' Model." *International Journal of Forecasting* 6:199–209.
- Bartolomei, S. M., and Sweet, A. L. (1989). "A Note on the Comparison of Exponential Smoothing Methods for Forecasting Seasonal Series." *International Journal of Forecasting* 5:111–116.
- Bhansali, R. J. (1980). "Autoregressive and Window Estimates of the Inverse Correlation Function." *Biometrika* 67:551–566.
- Bowerman, B. L., and O'Connell, R. T. (1979). *Time Series and Forecasting: An Applied Approach*. North Scituate, MA: Duxbury Press.
- Box, G. E. P., and Cox, D. R. (1964). "An Analysis of Transformations." *Journal of the Royal Statistical Society, Series B* 26:211–234.
- Box, G. E. P., and Jenkins, G. M. (1976). *Time Series Analysis: Forecasting and Control*. Rev. ed. San Francisco: Holden-Day.
- Box, G. E. P., and Tiao, G. C. (1975). "Intervention Analysis with Applications to Economic and Environmental Problems." *Journal of the American Statistical Association* 70:70–79.
- Brocklebank, J. C., and Dickey, D. A. (1986). *SAS System for Forecasting Time Series*. 1986 ed. Cary, NC: SAS Institute Inc.
- Brown, R. G. (1962). *Smoothing, Forecasting, and Prediction of Discrete Time Series*. New York: Prentice-Hall.
- Brown, R. G., and Meyer, R. F. (1961). "The Fundamental Theorem of Exponential Smoothing." *Operations Research* 9:673–685.

- Chatfield, C. (1978). "The Holt-Winters Forecasting Procedure." *Journal of the Royal Statistical Society, Series C* 27:264–279.
- Chatfield, C., and Prothero, D. L. (1973). "Box-Jenkins Seasonal Forecasting: Problems in a Case Study." *Journal of the Royal Statistical Society, Series A* 136:295–315.
- Chatfield, C., and Yar, M. (1988). "Holt-Winters Forecasting: Some Practical Issues." *The Statistician* 37:129–140.
- Chatfield, C., and Yar, M. (1991). "Prediction Intervals for Multiplicative Holt-Winters." *International Journal of Forecasting* 7:31–37.
- Cogger, K. O. (1974). "The Optimality of General-Order Exponential Smoothing." *Operations Research* 22:858–867.
- Cox, D. R. (1961). "Prediction by Exponentially Weighted Moving Averages and Related Methods." *Journal of the Royal Statistical Society, Series B* 23:414–422.
- Davidson, J. E. H. (1981). "Problems with the Estimation of Moving Average Processes." *Journal of Econometrics* 16:295–310.
- Dickey, D. A., and Fuller, W. A. (1979). "Distribution of the Estimators for Autoregressive Time Series with a Unit Root." *Journal of the American Statistical Association* 74:427–431.
- Dickey, D. A., Hasza, D. P., and Fuller, W. A. (1984). "Testing for Unit Roots in Seasonal Time Series." *Journal of the American Statistical Association* 79:355–367.
- Fair, R. C. (1986). "Evaluating the Predictive Accuracy of Models." In *Handbook of Econometrics*, vol. 3, edited by Z. Griliches and M. D. Intriligator, 1979–1995. New York: North-Holland.
- Fildes, R. (1979). "Quantitative Forecasting—the State of the Art: Extrapolative Models." *Journal of the Operational Research Society* 30:691–710.
- Fuller, W. A. (1976). *Introduction to Statistical Time Series*. New York: John Wiley & Sons.
- Gardner, E. S., Jr. (1984). "The Strange Case of the Lagging Forecasts." *Interfaces* 14:47–50.
- Gardner, E. S., Jr. (1985). "Exponential Smoothing: The State of the Art." *Journal of Forecasting* 4:1–38.
- Granger, C. W. J., and Newbold, P. (1977). *Forecasting Economic Time Series*. New York: Academic Press.
- Greene, W. H. (1993). *Econometric Analysis*. 2nd ed. New York: Macmillan.
- Hamilton, J. D. (1994). *Time Series Analysis*. Princeton, NJ: Princeton University Press.
- Harvey, A. C. (1981). *Time Series Models*. New York: John Wiley & Sons.
- Harvey, A. C. (1984). "A Unified View of Statistical Forecasting Procedures." *Journal of Forecasting* 3:245–275.
- Hopwood, W. S., McKeown, J. C., and Newbold, P. (1984). "Time Series Forecasting Models Involving Power Transformations." *Journal of Forecasting* 3:57–61.
- Jones, R. H. (1980). "Maximum Likelihood Fitting of ARMA Models to Time Series with Missing Observations." *Technometrics* 22:389–396.

- Judge, G. G., Griffiths, W. E., Hill, R. C., and Lee, T.-C. (1980). *The Theory and Practice of Econometrics*. New York: John Wiley & Sons.
- Ledolter, J., and Abraham, B. (1984). "Some Comments on the Initialization of Exponential Smoothing." *Journal of Forecasting* 3:79–84.
- Ljung, G. M., and Box, G. E. P. (1978). "On a Measure of Lack of Fit in Time Series Models." *Biometrika* 65:297–303.
- Makridakis, S. G., Wheelwright, S. C., and McGee, V. E. (1983). *Forecasting: Methods and Applications*. 2nd ed. New York: John Wiley & Sons.
- McKenzie, E. (1984). "General Exponential Smoothing and the Equivalent ARMA Process." *Journal of Forecasting* 3:333–344.
- McKenzie, E. (1986). "Error Analysis for Winters' Additive Seasonal Forecasting System." *International Journal of Forecasting* 2:373–382.
- Montgomery, D. C., and Johnson, L. A. (1976). *Forecasting and Time Series Analysis*. New York: McGraw-Hill.
- Morf, M., Sidhu, G. S., and Kailath, T. (1974). "Some New Algorithms for Recursive Estimation on Constant Linear Discrete Time Systems." *IEEE Transactions on Automatic Control* 19:315–323.
- Nelson, C. R. (1973). *Applied Time Series for Managerial Forecasting*. San Francisco: Holden-Day.
- Newbold, P. (1981). "Some Recent Developments in Time Series Analysis." *International Statistical Review* 49:53–66.
- Newton, H. J., and Pagano, M. (1983). "The Finite Memory Prediction of Covariance Stationary Time Series." *SIAM Journal on Scientific and Statistical Computing* 4:330–339.
- Pankratz, A. (1983). *Forecasting with Univariate Box-Jenkins Models: Concepts and Cases*. New York: John Wiley & Sons.
- Pankratz, A. (1991). *Forecasting with Dynamic Regression Models*. New York: John Wiley & Sons.
- Pankratz, A., and Dudley, U. (1987). "Forecast of Power-Transformed Series." *Journal of Forecasting* 6:239–248.
- Pearlman, J. G. (1980). "An Algorithm for the Exact Likelihood of a High-Order Autoregressive–Moving Average Process." *Biometrika* 67:232–233.
- Priestley, M. B. (1981). *Spectral Analysis and Time Series*. London: Academic Press.
- Roberts, S. A. (1982). "A General Class of Holt-Winters Type Forecasting Models." *Management Science* 28:808–820.
- Schwarz, G. (1978). "Estimating the Dimension of a Model." *Annals of Statistics* 6:461–464.
- Sweet, A. L. (1985). "Computing the Variance of the Forecast Error for the Holt-Winters Seasonal Models." *Journal of Forecasting* 4:235–243.
- Winters, P. R. (1960). "Forecasting Sales by Exponentially Weighted Moving Averages." *Management Science* 6:324–342.

- Woodfield, T. J. (1987). “Time Series Intervention Analysis Using SAS Software.” In *Proceedings of the Twelfth Annual SAS Users Group International Conference*, 331–339. Cary, NC: SAS Institute Inc.
<http://www.sascommunity.org/sugi/SUGI87/Sugi-12-57%20Woodfield.pdf>.
- Yar, M., and Chatfield, C. (1990). “Prediction Intervals for the Holt-Winters Forecasting Procedure.” *International Journal of Forecasting* 6:127–137.

Index

- additive-invertible region
 - smoothing weights, [4045](#)
- adjusted R-square
 - statistics of fit, [4062](#)
- adjustments, [4040](#)
- AIC, *see* Akaike's information criterion
- Akaike's information criterion
 - statistics of fit, [4062](#)
- Akaike's information criterion (AIC), [4062](#)
- Amemiya's prediction criterion
 - statistics of fit, [4062](#)
- Amemiya's R-square
 - statistics of fit, [4062](#)
- ARIMA model
 - autoregressive integrated moving-average model, [4053](#)
- autoregressive integrated moving-average model, *see* ARIMA model
- BIC, *see* Schwarz Bayesian information criterion
- boundaries
 - smoothing weights, [4045](#)
- Box-Cox transformation, *see* transformations
- Brown smoothing model, *see* double exponential smoothing
- calculations
 - smoothing models, [4042](#)
- combining forecasts, [4040](#)
- corrected sum of squares
 - statistics of fit, [4061](#)
- cubic trend, [4058](#)
- cubic trend curves, *see* trend curves
- damped-trend exponential smoothing, [4049](#)
 - smoothing models, [4049](#)
- diagnostic tests, [4060](#)
- Dickey-Fuller test, [4061](#)
- double exponential smoothing, [4047](#)
 - Brown smoothing model, [4047](#)
 - smoothing models, [4047](#)
- error sum of squares
 - statistics of fit, [4062](#)
- exponential smoothing, *see* smoothing models
- exponential trend, [4058](#)
- exponential trend curves, *see* trend curves
- external forecasts, [4040](#)
- forecast combination, [4040](#)
- forecasting, [4038](#)
- forecasting models
 - smoothing models, [4042](#)
 - transfer functions, [4055](#)
- forecasts
 - external, [4040](#)
- goodness-of-fit statistics, *see* statistics of fit
- Holt smoothing model, *see* linear exponential smoothing
- Holt-Winters method, *see* Winters method
- hyperbolic trend, [4058](#)
- hyperbolic trend curves, *see* trend curves
- initializations
 - smoothing models, [4043](#)
- intervention notation, [4059](#)
- interventions, [4058](#)
 - point, [4058](#)
 - predictor variables, [4058](#)
 - ramp, [4059](#)
 - step, [4059](#)
- linear exponential smoothing, [4048](#)
 - Holt smoothing model, [4048](#)
 - smoothing models, [4048](#)
- linear trend, [4058](#)
- linear trend curves, *see* trend curves
- log test, [4060](#)
- log transformation, *see* transformations
- logarithmic trend, [4058](#)
- logarithmic trend curves, *see* trend curves
- logistic transformation, *see* transformations
- logistic trend, [4058](#)
- logistic trend curves, *see* trend curves
- mean absolute error
 - statistics of fit, [4062](#)
- mean absolute percent error
 - statistics of fit, [4062](#)
- mean percent error
 - statistics of fit, [4063](#)
- mean prediction error
 - statistics of fit, [4062](#)
- mean squared error
 - statistics of fit, [4062](#)
- missing values

- smoothing models, 4044
- MMAE, 4042
- MMSE, 4042
- model evaluation, 4036
- multiplicative seasonal smoothing, 4051
 - smoothing models, 4051
- nonmissing observations
 - statistics of fit, 4061
- nonseasonal ARIMA model
 - notation, 4054
- nonseasonal transfer function
 - notation, 4055
- notation
 - nonseasonal ARIMA model, 4054
 - nonseasonal transfer function, 4055
 - seasonal ARIMA model, 4054
 - seasonal transfer function, 4056
- number of observations
 - statistics of fit, 4061
- optimizations
 - smoothing weights, 4045
- parameter estimation, 4036
- point interventions, 4058
- power curve trend, 4058
- power trend curves, *see* trend curves
- predictions
 - smoothing models, 4044
- predictor variables
 - interventions, 4058
 - seasonal dummies, 4060
 - trend curves, 4057
- quadratic trend, 4058
- quadratic trend curves, *see* trend curves
- R-square statistic
 - statistics of fit, 4062
- ramp function, *see* ramp interventions
- ramp interventions, 4059
 - ramp function, 4058
- random walk R-square
 - statistics of fit, 4062
- root mean square error
 - statistics of fit, 4062
- SBC, *see* Schwarz Bayesian information criterion
- Schwarz Bayesian information criterion
 - BIC, 4062
 - SBC, 4062
 - statistics of fit, 4062
- seasonal ARIMA model
 - notation, 4054
- seasonal dummies, 4060
 - predictor variables, 4060
- seasonal exponential smoothing, 4050
 - smoothing models, 4050
- seasonal transfer function
 - notation, 4056
- seasonality test, 4061
- series adjustments, 4040
- series diagnostics, 4060
- simple exponential smoothing, 4046
 - smoothing models, 4046
- smoothing equations, 4043
 - smoothing models, 4043
- smoothing models
 - calculations, 4042
 - damped-trend exponential smoothing, 4049
 - double exponential smoothing, 4047
 - exponential smoothing, 4042
 - forecasting models, 4042
 - initializations, 4043
 - linear exponential smoothing, 4048
 - missing values, 4044
 - multiplicative seasonal smoothing, 4051
 - predictions, 4044
 - seasonal exponential smoothing, 4050
 - simple exponential smoothing, 4046
 - smoothing equations, 4043
 - smoothing state, 4043
 - smoothing weights, 4044
 - standard errors, 4045
 - underlying model, 4042
 - Winters method, 4051, 4053
- smoothing state, 4043
 - smoothing models, 4043
- smoothing weights, 4044
 - additive-invertible region, 4045
 - boundaries, 4045
 - optimizations, 4045
 - smoothing models, 4044
 - specifications, 4045
 - weights, 4044
- specifications
 - smoothing weights, 4045
- square root transformation, *see* transformations
- standard errors
 - smoothing models, 4045
- statistics of fit, 4061
 - adjusted R-square, 4062
 - Akaike's information criterion, 4062
 - Amemiya's prediction criterion, 4062
 - Amemiya's R-square, 4062
 - corrected sum of squares, 4061
 - error sum of squares, 4062
 - goodness-of-fit statistics, 4061

- mean absolute error, [4062](#)
- mean absolute percent error, [4062](#)
- mean percent error, [4063](#)
- mean prediction error, [4062](#)
- mean squared error, [4062](#)
- nonmissing observations, [4061](#)
- number of observations, [4061](#)
- R-square statistic, [4062](#)
- random walk R-square, [4062](#)
- root mean square error, [4062](#)
- Schwarz Bayesian information criterion, [4062](#)
- uncorrected sum of squares, [4061](#)
- step function, *see* step interventions
- step interventions, [4059](#)
 - step function, [4058](#)
- transfer functions, [4055](#)
 - forecasting models, [4055](#)
- transformations
 - Box-Cox, [4041](#)
 - log, [4041](#)
 - logistic, [4041](#)
 - square root, [4041](#)
- trend curves, [4057](#)
 - cubic, [4058](#)
 - exponential, [4058](#)
 - hyperbolic, [4058](#)
 - linear, [4058](#)
 - logarithmic, [4058](#)
 - logistic, [4058](#)
 - power, [4058](#)
 - predictor variables, [4057](#)
 - quadratic, [4058](#)
- trend test, [4061](#)
- uncorrected sum of squares
 - statistics of fit, [4061](#)
- underlying model
 - smoothing models, [4042](#)
- weights, *see* smoothing weights
- Winters method, [4051](#), [4053](#)
 - Holt-Winters method, [4051](#)
 - smoothing models, [4051](#), [4053](#)